


Frances Cabaniss Roberts Collection: Series 10, Box 6, Folder 3  
Orton, James "Underground Treasures: How and Where to Find Them"

Image 1 r10\_06-03-000-0001 [Contents](#) [Index](#) [About](#)




**Names:**

Underground  
Treasures

**Types:**

book cover



**Names:**

Snodgrass, John


**Places:**

Scottsboro, AL

**Types:**

signatur

Frances Cabaniss Roberts Collection: Series 10, Box 6, Folder 3  
Orton, James "Underground Treasures: How and Where to Find Them"  
Image 3 r10\_06-03-000-0003 [Contents](#) [Index](#) [About](#)



**Names:**

Orton, James

Underground  
Treasures

**Places:**


Hartford, CT

**Types:**

book



Frances Cabaniss Roberts Collection: Series 10, Box 6, Folder 3  
Orton, James "Underground Treasures: How and Where to Find Them"  
Image 4 r10\_06-03-000-0004 [Contents](#) [Index](#) [About](#)



**Names:**

Orton, James

Underground  
Treasures

**Places:**

Hartford, CT


**Types:**

book

**Dates:**

1872

Frances Cabaniss Roberts Collection: Series 10, Box 6, Folder 3  
Orton, James "Underground Treasures: How and Where to Find Them"  
Image 5 r10\_06-03-000-0005 [Contents](#) [Index](#) [About](#)



**Names:**


Raymond, Rossiter  
W.

**Types:**

book

**Dates:**

1872




**Names:**

Underground  
Treasures

**Types:**

book

note



**Names:**


Illustrations

Table of Contents

**Types:**

list






**Names:**


Chapter I - Table of Contents  
 Introduction


**Types:**

book list









**Names:**

Chapter II -


Key

Directions - The

**Types:**

book







Series 10, Box 6, Folder 3

18 UNDERGROUND TREASURES.

---

**A.**

WILL SCRATCH GLASS.

I. OPAQUE.

(1) Black: 12, 20, 28, 30, 42, 43, 51, 54, 67, 72.  
 (2) Brown: 12, 28, 42, 59, 72, 77.  
 (3) Red: 39, 46, 54, 59, 67.  
 (4) Yellow: 38, 72, 77.  
 (5) Gray: 22, 28, 72.  
 (6) White: 64.

II. TRANSLUCENT.

(1) Brown: 32, 39, 72, 77.  
 (2) Red: 17, 32, 46, 59, 68, 73, 74.  
 (3) Yellow: 37, 39, 72, 73, 78.  
 (4) Green: 74, 77.  
 (5) Violet blue: 3.  
 (6) White: 27, 57, 73, 77.  
 (7) Banded or clouded: 2.

19

THE KEY.

---

**B.**

WILL NOT SCRATCH GLASS.


I. OPAQUE.

|   |  |
|---|--|
| <p style="text-align: center;">Harder than white marble.*</p> <p>(1) Black: 12, 35, 47.<br/>                 (2) Brown: 66, 75.<br/>                 (3) Red: 44, 53, 75.<br/>                 (4) Yellow: 26, 44.<br/>                 (5) Green: 45.<br/>                 (6) Gray: 25, 66.<br/>                 (7) White: 6, 9, 12.</p> | <p style="text-align: center;">Softer than white marble.</p> <p>4, 7, 10, 12, 13, 14, 16, 24,<br/>                 34, 37, 49, 51, 55, 56, 76,<br/>                 12, 14, 21.<br/>                 21, 23, 41, 55.<br/>                 22, 33, 56.<br/>                 60, 62.<br/>                 5, 22, 32, 34, 36, 43, 56, 63,<br/>                 69.<br/>                 36, 42, 58, 62.</p> |
|---|--|

II. TRANSLUCENT.

|   |   |
|---|---|
| <p>(1) Black: 11.<br/>                 (2) Brown: 9, 11, 65, 66.<br/>                 (3) Red: 9, 11, 18, 53, 78.<br/>                 (4) Yellow: 9, 11, 15, 29,<br/>                 47, 78.<br/>                 (5) Green: 29, 45, 65, 70.<br/>                 (6) Blue: 8, 18, 29, 47.<br/>                 (7) Gray: 19, 47, 65, 66.<br/>                 (8) White: 18, 47.<br/>                 (9) Mottled or Banded: 47.</p> | <p>55.<br/>                 48.<br/>                 36, 55.<br/>                 48, 71.<br/>                 48, 60, 61.<br/>                 57, 69.<br/>                 2.</p> |
|---|---|

\*That is, they are not so easily cut with a knife; they do not necessarily scratch marble.



**Names:**

Chapter III -  
Descriptive List of

Minerals

**Types:**

book

streaks. Cape Sable, Md.; Cleveland County, N. C.; coal slates on Ohio River, and in caves in Sevier, De Kalb, Coffee and Franklin Counties, Tenn.; also Esmeralda and Storey Counties, Nev.

3—AMETHYST.

Same as *Rock Crystal*, but colored purple or bluish violet. Generally in clustered crystals.

VALUE.—When clear and finely colored, it is a favorite gem.

LOCALITIES.—Usually found with agate. Keweenaw Point, Pic Bay and Gargontwa on Lake Superior; Bristol, R. I.; Surry, N. H.; East Bradford, Aston, Chester, Thorabury, Edgemont, Sadsbury, Birmingham, Middletown and Providence, Penn.; Greensboro, N. C.

4—ANTHRACITE.

Occurs massive; compact; high lustre; brittle; breaks with a curved surface; will not scratch marble; burns, but not readily, with a pale blue flame and little smoke;

will not form coke by roasting; gravity 1.4 to 1.8.

VALUE.—Used for fuel and sometimes cut into inkstands, etc.

LOCALITIES.—Found in beds between slates and sandstones, and east of the Alleghany range only, as Eastern Pennsylvania; Portsmouth, R. I.; Mansfield, Mass.; North Carolina. No workable beds will be found in New York.

The rocks in anthracite regions are tilted, bent and broken, never level to any great extent. Impressions of leaves are good indications.

5—ANTIMONY ORE.


Occurs fibrous or granular; color lead gray, often tarnished; shining lustre, brittle; but thin pieces can be cut off with a knife; melts in a candle, at a high heat passing off in vapor; gravity 4.5.

VALUE.—The source of the antimony of commerce, containing seventy per cent.

LOCALITIES.—Found associated with *Silver, Spathic Iron, Blende, Baryta* and

Series 10, Box 6, Folder 3





*Quartz.* Carmel, Me.; Lyme, N. H.; Soldier's Delight, Md.; Aurora, Nev.; San Amedeo Cañon and Tulare County, Cal.

6—ASBESTUS.

Occurs finely fibrous, flax-like; flexible, not elastic; silky lustre, sometimes greenish; gravity 3.

VALUE.—Used for lining safes and steam-packing, and for making incombustible cloth, lamp-wicks, etc.

LOCALITIES.—Found in granite-regions east of the Alleghanies; often with *Serpentine*. Brighton, Dedham, Newbury, Pelham and Sheffield, Mass.; Milford, West Farms, Winchester and Wilton, Conn.; Chester, Mt. Holly and Cavendish, Vt.; Patterson, Philipstown, Monroe and Staten Island, N. Y.; Brunswick, N. J.; East Nottingham, Gosben and Aston, Penn.; Bare Hills and Cooptown, Md.; Barnett's Mills, Va.

7—ASPHALTUM.

Occurs massive; brittle; breaking with high lustre like hardened tar, and with

Series 10, Box 6, Folder 3

curved surface; melts and burns readily with flame and smoke; gravity 1.2, sometimes floats on water.

VALUE.—Used for cements and varnishes.

LOCALITIES.—Found generally near the surface. Near the coast of Santa Barbara, Cal.; West Virginia, twenty miles south of Parkersburg.

8—AZURITE.

Occurs in crystals and masses with glassy lustre, or earthy and dull; brittle; crackles and blackens, and finally fuses by heat; dissolves with effervescence in nitric acid; gravity 3.5.

VALUE.—A valuable ore of copper, containing sixty per cent.

LOCALITIES.—Found chiefly in lead and copper mines. Perkiomen lead mine, Cornwall, Phoenixville and Nicholson's Gap, Pa.; near New Brunswick, N. J.; near Mineral Point, Wis.; Polk County, Tenn.; Calaveras and Mariposa Counties, Cal.; near Virginia City, Mont.



Series 10, Box 6, folder 3

9.—BARTTA, OR HEAVY SPAR.

Occurs in crystals, plates and masses; powder white; brittle; crackles when strongly heated; not dissolved in acids; easily distinguished by its weight; gravity 4.5, or twice as heavy as *Gypsum*.

VALUE.—Used extensively as white paint and in pottery.

LOCALITIES.—Found in mining districts, often with lead, copper and iron ores, and in limestone. Piermont, N. H.; Hatfield, Southampton and Leverett, Mass.; Cheshire and Berlin, Conn.; Pillar Point, Rossie, Carlisle, Schoharie, De Kalb, Gouverneur, N. Y.; Fauquier and Buckingham Counties, Va.; Union, Gaston and Orange Counties, N. C.; near Paris, and in Anderson, Fayette, Mercer and Owen Counties, Ky.; on Brown's Creek and Haysboro, Tenn.; Bainbridge, O.; Scales Mound, Ill.; Prince Vein, Lake Superior; Mine-a-Barton, Mo.; near Fort Wallace, N. M.; Ingo County, Cal.

10.—BITUMINOUS COAL.

Occurs in masses, beds or seams; softer and duller than *Anthracite*; often a bright pitchy lustre; brittle, showing a slaty or jointed structure rather than curved surface; powder black; burns readily with yellow flame; by roasting forms coke; gravity 1.5 or less.

VALUE.—Used for fuel and the production of gas, coke, carbolic acid and aniline.

LOCALITIES.—Found west of Harrisburg, Pa., in rocks (slates and sandstones) less disturbed than in the *Anthracite* region. Western Pennsylvania; South-east Ohio; West Virginia; Eastern Kentucky and Tennessee to Tuscaloosa; North-west Kentucky; Illinois; Iowa; Missouri; Kansas; Arkansas; Northern Texas; Central Michigan; Owyhee County, Idaho; Deer Lodge and Gallatin Counties and sixty miles north-east of Bannock, Mont.

11.—BLENDE.

Occurs in crystals and masses; waxy lustre, but not always very apparent; usual

color, rosin-yellow to dark brown; brittle; the powder, which is whitish to reddish-brown, dissolves in muriatic acid giving off the odor of rotten eggs; by roasting gives off sulphur-fumes; infusible alone, but on charcoal at a high heat gives off white fumes; gravity 4.

VALUE.—An ore of zinc (containing sixty-six per cent.) and a source of white vitriol. Often worked for its *Silver* and *Gold*.

LOCALITIES.—Found with lead and other ores. Lubec and Bingham, Me.; Eaton, Warren and Shelburne, N. H.; Sterling, Southampton and Hatfield, Mass.; Brookfield, Berlin, Roxbury and Monroe, Conn.; near Wurtsboro, Cooper's Falls, Mineral Point, Fowler, Ancram, Clinton and Spraker's Basin, N. Y.; Wheatley and Perkiomen lead-mines, Schoeykill, Shannonville and Friedensville, Pa.; Austin's lead-mine, Va.; Haysboro', Brown's Creek and Polk Counties, Tenn.; Prince Vein, Mich.; Duboquet, Ia.; Warsaw, Rosiclare and Ga-


lena, Ill.; Shullsburg, Wis.; Stillwater, Minn.

12.—BOG IRON ORE.

Occurs in masses or beds, looking much like hard brown earth; loose or porous and earthy, rather than compact and nodular; powder yellowish-brown; when strongly heated becomes black and magnetic; gravity nearly 4. An earthy yellow variety is called *Yellow Oxide*.

VALUE.—An important ore, yielding thirty-five per cent.

LOCALITIES.—Found in low, marshy grounds; widely distributed Lebanon, N. H.; Berkshire and Plymouth Counties, Mass.; Columbia, St. Lawrence, Franklin and Jefferson Counties, N. Y.; New Limerick, Katahdin, Newfield, Shapleigh, Argyle, Clinton, Williamsburg and Lebanon, Me.; Darien and Martin Counties, Ind.; Monmouth County, N. J.; Somerset and Worcester Counties, Md.; Michigan, Ohio, Illinois, Wisconsin, etc.



13.—BRITTLE SILVER ORE.

Occurs in crystals and masses; metallic lustre; tarnishes yellow, gray and finally black; easily cut or broken; when heated gives off fumes of sulphur and antimony, affording a button of silver; dissolved in nitric acid, it silvers copper placed in it; gravity 6.

VALUE.—A rich ore of silver, containing over sixty per cent.

LOCALITIES.—Found in veins with other silver ores, in Nevada and Idaho.

14.—BROWN COAL.

Occurs like *Bituminous Coal*, but usually brownish-black with less lustre, and often showing a woody or slaty structure; powder always brown; contains fossil plants; gravity between 1.2 and 1.5.

VALUE.—Inferior to No. 10. Makes no coke. Can be used in the manufacture of alum.

LOCALITIES.—Found in thin veins or elliptical masses, never in extensive layers like Pennsylvania coal. Near Richmond,

Va.; Deep River, N. C.; Michigan, Missouri, Texas; Evanston, Utah; Coal Creek and Bellmonte, Col.; Boreman, Dearborn River and Greenhorn Gulch, Mont.

15.—CALAMINE.

Occurs in crystals and masses; glossy lustre; harder than marble; brittle; heated it swells up, becomes opaque and emits a green light; dissolves, when powdered, in hot sulphuric acid without effervescence; gravity 3.4.

VALUE.—An ore of zinc yielding from forty to sixty per cent.

LOCALITIES.—Found in limestone rock with other ores. Friedensville, Perkiomen, Phoenixville, Lancaster and Selin's Grove, Pa.; Austin's Mines in Wythe County, Va.; Claiborne County, Tenn.; Jefferson County, Mo.

16.—CANNEL COAL.

Occurs in compact masses; dull lustre; brittle, breaking with a curved surface;



Series 10, Box 6, Folder 3

burns readily but does not melt; does not soil the fingers; gravity about 1.2.

VALUE.—Used for fuel and for making gas, oil and ornaments.

LOCALITIES.—Found in the Mississippi Valley; Kentucky; Lick, Ohio; Illinois; Moniteau County, Mo.; Kenawha County, Va.; Beaver County, Pa.

17.—CARNELIAN.

Occurs in masses or pebbles; at first grayish, but by exposure to the sun becomes uniform flesh, red or brown, never striped,—although *Carnelian* may form one of the bands of an *Agate*; brittle, breaking with a curved surface; very hard; takes a fine polish; glassy or resinous lustre; gravity 2.6.

VALUE.—Used for jewelry. When of two layers, white and red, (properly called *sardonyx*.) it is used for cameos.

LOCALITIES.—Same as *Agate*.

18.—CELESTINE.

Occurs crystallized, fibrous and massive; color white, often faint bluish; glassy lus-

tre; very brittle; under the blow-pipe crackles and melts, tinging the flame red; does not dissolve in acids; gravity 4.

VALUE.—The source of nitrate of strontia, used in fire-works.

LOCALITIES.—Found in limestone, gypsum and sandstone. Rossie, Schoharie, Chaumont Bay, Depauville and Stark, N. Y.; Frankstown, Pa.; Strontian and Put-in-Bay Islands, Lake Erie; near Nashville, Tenn.; Fort Dodge, Iowa.


19.—CERUSSITE.

Occurs in crystals, in powder or masses; glassy lustre; brittle; dissolves in nitric acid with effervescence; heated strongly on charcoal crackles and fuses, giving a globule of lead; gravity 6.4.

VALUE.—A rich ore of lead yielding seventy-five per cent.

LOCALITIES.—Found in lead mines. Southampton, Mass.; Perkiomen, Phoenixville, Charlestown and Schuylkill, Pa.; Wythe County, Va.; Washington Mine, N. C.; Valle's Diggings, Mine-la-Motte





and Mine-a-Burton, Mo.; Davies and Rock Counties, Ill.; Blue Mounds, Wis.; Ingo County, Cal.

20.—CHROMIC IRON.

Occurs in compact masses; powder dark brown; small pieces sometimes attracted by the magnet; brittle, breaking with uneven surface; with borax melts into a green globule; not acted upon by acids; little lustre; gravity 4.4.

VALUE.—Used in making the chrome pigments.

LOCALITIES.—Found in *Serpentine*. Bare Hills, Cooptown and north part of Cecil County, Md.; Nottingham, W. Gosben, Williston, Fulton, Mineral Hill, Texas and Unionville, Pa.; Jay, New Fane, Westfield and Troy, Vt.; Chester and Blanford, Mass.; Loudon County, Va.; Yancy County, N. C.; North Almaden, New Idria and Coloma, Cal.

21.—CINNABAR.

Occurs in granular or earthy masses; resembles iron-rust, but is a yellowish-red;

powder scarlet; easily cut with a knife; thrown on red-hot iron, evaporates, giving off odor of sulphur; rubbed on copper, "silvers" it; gravity 9, or about as heavy as *Copper*.

VALUE.—The source of mercury (containing eighty-four per cent.) and vermilion.

LOCALITIES.—Found in slate and limestone rocks. Centreville, Coulterville, New Idria and New Almaden, and Lake and San Luis Obispo Counties, California; Idaho.

22.—COBALT PYRITES.

Occurs crystallized and massive; does not scratch glass easily; metallic lustre; tarnish, copper-red; powder, blackish-gray; brittle; heated on charcoal gives off sulphur fumes; heated with borax gives a blue glass; gravity 5.

VALUE.—An ore of cobalt, yielding twenty per cent.

LOCALITIES.—Usually found in slate or granite rocks with *Copper Pyrites*. Mineral Hill, Md.; Mine-la-Motte, Mo.

Series 10, Box 6, Folder 3

23.—COPPER.

Occurs in irregular masses; metallic lustre; can be cut with a knife; malleable; ductile; fusible; gravity 8.8.

VALUE.—A source of copper and silver.

LOCALITIES.—Most abundant in the trap and "freestone" regions. New Brunswick, Somerville, Schuyler's and Flemington, N. J.; Whately, Mass.; Cornwall and Shannonville, Pa.; Polk County, Tenn.; Keweenaw Point, Lake Superior; Calaveras, Amador and Santa Barbara Counties, Cal.; on Gila River, Ariz.

24.—COPPER GLANCE.

Occurs crystallized and massive; color, blackish lead-gray, often tarnished blue or green; nearly as hard as marble; brittle; a splinter will melt in a candle, giving off the odor of sulphur; dissolved in nitric acid, it will coat a knife-blade with copper; metallic lustre; gravity 5.5.

VALUE.—An ore of copper, yielding seventy-five per cent.

LOCALITIES.—Found at copper-mines.

Simsbury, Bristol and Cheshire, Conn.; Schuyler's Mines, N. J.; Orange County, Va.; near Newmarket, Md.; Lake Superior copper-region; La Paz, Arizona; Washoe, Humboldt, Nye and Churchill Counties, Nev.

25.—COPPER NICKEL.


Occurs in masses; metallic lustre; color pale copper-red; tarnishes gray to black; powder pale brownish-black; brittle; on charcoal melts giving the odor of garlic; becomes green in nitric acid; gravity 7.5.

VALUE.—An ore of nickel (containing forty-four per cent.) and arsenic.

LOCALITIES.—Found in granite regions. Chatham, Conn.

26.—COPPER PYRITES.

Occurs in crystals and masses; color brass-yellow; tarnishes green; metallic lustre when freshly broken; can be cut with a knife; brittle; powder greenish black; on charcoal melts giving off sulphur fumes; dissolves in nitric acid, making a green liquid; gravity 4.2.




VALUE.—If of a fine yellow hue, it is a valuable copper ore (yielding from twelve to forty per cent.) and source of blue vitriol.

LOCALITIES.—Found in mountainous or granite regions with other ores. Lubec and Dexter, Me.; Franconia, Unity, Warren, Eaton, Lyme, Haverhill and Shelburne, N. H.; Corinth, Waterbury and Strafford, Vt.; Southampton, Turner's Falls, Hatfield and Sterling, Mass.; Bristol and Middletown, Conn.; Ancram, Rossie, Wurtzboro' and Ellenville, N. Y.; Phoenixville and Pottstown, Pa.; Bare Hills, Catoctin Mountains, near Newmarket and Finksbury, Md.; Phoenix and Walton Mines, Va.; Greensboro, Charlotte and Phoenix Mines, N. C.; Hiwassee Mines, Tenn.; Cherokee, Rabun and Habersham Counties, Ga.; Presque Island, Lake Superior; Mineral Point, Wis.; Union, Keystone, Empire and other mines, Calaveras County, La Victoire and Haskell claims in Mariposa County, Amador and Plumas Counties, Cal.; near Virginia City, Mont.



Series 10, Box 6, folder 3



PERSPECTIVE DIAMOND DRILL.

DIAMOND—EMERY. 41

27.—DIAMOND.

Occurs in crystals and irregular angular masses; cannot be scratched by any other mineral or the file; brilliant lustre; feels cold to the touch; when rubbed on the sleeve exhibits electricity for hours; retains the breath but a short time; often tinged yellow, red, or green; gravity 3.5.

VALUE.—Used for jewelry, lenses and for cutting glass.

LOCALITIES.—Found in gold-regions, in river-washings of sand and pebbles; usually with coarse gold, but deeper down. Rutherford, Cabarras, Franklin and Lincoln Counties, N. C.; Hall County, Ga.; Manchester, Va.; Cherokee Ravine, N. San Juan, French Canal, Forrest Hill, Placerville and Fiddletown, Cal.

28.—EMERY.

Occurs in granular masses, sometimes with bluish crystals; looks like fine grained iron ore; breaks with uneven surface; scratches quartz easily; very tough; brittle; gravity 4.

VALUE.—Used extensively as a cutting and polishing material.

LOCALITIES.—Found generally in limestone or granite with *Magnetic Iron Ore*. Chester, Mass.; Newlin and Unionville, Penn.; Macon and Guilford Counties, N. C.

29.—FLUOR SPAR.

Occurs in square crystals and in masses; glassy lustre; powder white; brittle; crackles when heated and then shines in the dark; does not effervesce with acids; is not scratched by marble; gravity 3.

VALUE.—Used as flux in glass and iron works.

LOCALITIES.—Found in limestone, granite, slate, etc., often at lead-mines. Blue Hill Bay, Me.; Westmoreland, N. H.; Putney, Vt.; Southampton, Mass.; Trumbull, Plymouth, Middletown and Willimantic, Conn.; Muscolonge Lake, Rossie and Johnsburg, N. Y.; near Franklin, N. J.; near Woodstock and Shepardstown, Va.; Smith County, Tenn.; Mercer County,

Series 10, Box 6, folder 3

Ky.; Gallatin County, along the Ohio, Ill. Castle Dome District, Ariz.

30.—FRANKLINITE.

Occurs crystallized and in masses; generally made of coarse grains; brittle; powder dark reddish-brown; heated with soda turns bluish-green; dissolves in muriatic acid; gravity 5.

VALUE.—An ore of zinc.

LOCALITIES.—Found in limestone with *Garnet and Zincolite*. Hamburg and Stirling Hill, N. J.

31.—GALENA.

Occurs in crystals and masses; brilliant lustre; brittle; easily broken; powder, when finely rubbed is black; can be cut with a knife; heated it gives off sulphur and melts; dissolves in nitric acid leaving a white powder at the bottom; gravity 7.5—or a little heavier than cast-iron.

VALUE.—The main source of lead (yielding eighty per cent), and also smelted for the silver it contains. Used also in glazing stone-ware.

Series 10, Box 6, folder 3

LOCALITIES.—Generally found in limestone with *Iron Pyrites*, zinc-ore, etc. That found in slate is richest in silver. Abounds in Missouri, Illinois, Iowa, Wisconsin and Arkansas; Rossie, Wurtzboro, Ancram, Macomb and Ellenville, N. Y.; Lubec, Blue Hill Bay, Bingham and Parsonsville, Me.; Eaton, Shelburne, Haverhill, Warren and Bath, N. H.; Thetford, Vt.; Southampton, Leverett and Sterling, Mass.; Middletown and Roxbury, Conn.; Phoenixville, Charlestown, Schuylkill, Pequea Valley and Shannonville, Pa.; Austin's and Walton's Mines, Va.; Cabarras County, N. C.; Brown's Creek and Haysboro, Tenn.; Chocolate River, Mich.; Ingo County, Cal.; on Walker's River and Steamboat Springs, Nev.; Castle Dome and Eureka, Ariz.; Clear Creek County, Col.; Virginia City and Red Bluff Lode, Mont.; Cache Valley, Utah.

32.—GARNET.

Occurs in crystals with four-sided faces; often nearly round; deep red, which grows

darker by heat; rarely yellow; also in brown masses; melts at a high heat; brittle; not scratched by a knife; glassy lustre; gravity 4.

VALUE.—The clear deep red and yellow varieties are used for jewelry; the massive brown is ground for "emery."

LOCALITIES.—Found in slate and granite rocks. Bethel, Parsonsfield, Phippsburg, Windham, Brunswick and Ranford, Me.; Hanover, Franconia, Haverhill, Warren, Unity, Lisbon and Grafton, N. H.; New Fane, Cabot and Cavendish, Vt.; Carlisle, Boxborough, Brookfield, Brimfield, Newbury, Bedford, Chesterfield and Barre, Mass.; Reading, Monroe, Haddam and Middletown, Conn.; Rogers' Rock, Crown Point, Willsboro, Middletown, Amity, and near Yonkers, N. Y.; Franklin, N. J.; Pennsbury, Warwick, Aston, Knauertown, Chester, Leiperville and Mineral Hill, Pa.; Dickson's Quarry, Del.; Hope Valley, Cal.; near Virginia City, on Yellowstone and Madison Rivers, Mont.



33.—GOLD.

Occurs in scales, grains and nuggets, or disseminated through cellular quartz; metallic lustre; without tarnish; can be cut and hammered into thin plates; not dissolved by nitric acid; gravity 19, when pure and of a rich gold yellow color. The pale or brass yellow specimens are much lighter, the gravity being as low as 13. A grayish yellow gold, occurring in small, flat grains has a gravity of about 16.

LOCALITIES.—Found in veins of quartz running through greenish or grayish slates, the quartz at the surface being generally full of cavities and rusted, and the slates below the surface often containing little cubic crystals of *Iron Pyrites*: also in the valleys traversed by mountain-streams and in the river sands and gravel below. *Iron* and *Copper Pyrites*, *Galena* and *Blende* frequently contain gold. Masses of quartz and pyrites from the gold-regions, which make no show of gold, sometimes pay well; the value of such specimens can be




Series 10, Box 6, folder 3


Series 10, Box 6, Folder 3

THE MINERALS TREATED. 49

determined only by an assayer. Eastern range of Appalachians, as Habersham, Rabun, Clark, Hall, Lumpkin and Lincoln Counties, Ga.; Abbeville, Chesterfield, Union, Lancaster and Pickens Counties, S. C.; Montgomery, Cabarras, Mecklenburg, Burke and Lincoln Counties, N. C.; Spotsylvania, Buckingham, Fauquier, Stafford, Culpepper, Orange, Goochland and Louisa Counties, Va.; Dedham, Mass.; Bridgewater, Vt.; Canaan and Lisbon, N. H.; on Sandy River and Madrid, Me. Numberless points along the higher Rocky Mountains and western slope of Sierra Nevada, as near Santa Fe, Cerillos and Avo, New Mex.; San Francisco, Wauba and Yuma District, Ariz.; between Long's Peak and Pike's Peak, Col.; Comstock Lode, Nev.; Owyhee, Boise and Flint Districts and Poorman Lode, Idaho; Emigrant and Alder Gulches, Red Bluff and near Jefferson River, Mont.; Josephine District, Powder, Burnt, and John Day Rivers, western slope of Cascade Mountains, and







## 36.—GYPSUM.

Occurs in plates, fibres coarse and fine, and massive; pearly or glistening; powder white, which if heated and mixed with water, turns hard; does not dissolve in sulphuric acid; may be scratched by the nail; gravity 2.3.

VALUE.—Used for stucco, manure, glazing, statuary, manufacture of glass, etc. A variety, called *Satin Spar*, worked into necklace beads and other ornaments, is finely fibrous and compact, taking a polish (though easily scratched,) and then resembles pearl or opal.

LOCALITIES.—Found with marl or clay, limestone and salt. Camillus, Manlius, Stark and Lockport, N. Y.; on the St. Mary's and Patuxent, Md.; Washington County and Lynchburg, Va.; Charleston, S. C.; Poland, Ottawa and Canfield, O.; Davidson and Summer Counties, Tenn.; Grand Rapids and Sagenaw Bay, Mich.; Des Moines River, Iowa; Walker Lake and Six Mile Cañon, Nev.; Fort Dodge.

## 37.—HORN SILVER.

Occurs in crystals, wax-like masses, or in crusts; when scratched shows a shining streak; becomes brown on exposure; quite soft, easily cut; a small piece placed on zinc and moistened, swells up, turns black and shows metallic silver on being pressed with a knife; dissolves in harts-horn; gravity 5.5.


VALUE.—An ore of silver, yielding seventy per cent.

LOCALITIES.—Found in slate with other silver ores. Lake Superior Mining Region; Austin and Comstock Lode, Nev.; Willow Springs and San Francisco districts, Eldorado Cañon, Ariz.; Poorman Mine, Idaho.

## 38.—IRON PYRITES.

Occurs in masses and square crystals; splendid lustre; color, bronze-yellow; brittle; strikes fire with steel; heated it gives off sulphur fumes; powder brownish; gravity 5.

VALUE.—Affords sulphur, copperas and



alum. When found outside of the coal region, it often contains gold and silver.

LOCALITIES.—Found in all kinds of rocks. Bingham, Corinna, Farmington, Waterville, Brooksville, Peru and Jewett's Island, Me.; Shelburne, Unity and Warren, N. H.; Baltimore, Hartford and Shoreham, Vt.; Heath, Hubbardston and Hawley, Mass.; Roxbury, Monroe, Orange, Milford, Middletown, Stafford, Colchester, Ashford, Tolland and Union, Conn.; Rossie, Malone, Phillips, Johnsburgh, Canton, Chester, Warwick and Franklin, Putnam and Orange Counties, N. Y.; Chester, Knauertown, Cornwall and Pottstown, Pa.; Greensboro', N. C.; Mercer County, Ky.; Bainbridge, O.; Galema at Marsden's Diggings, Ill.; on Sugar Creek, Ind.; mines of Colorado and California.

39.—JASPER.

Occurs in masses, either in veins or as rounded stones; dull lustre, yet takes a high polish; breaks with a curved surface;

not attacked by acids; is scratched by *Rock Crystal*; gravity 2.5.

VALUE.—Used for mosaics and other ornaments when compact, fine-grained and bright color.

LOCALITIES.—Found everywhere. Sugar Loaf Mountain and Machiasport, Me.; Saugus, Mass.; Castleton and Colchester, Vt.; Bloomingrove, N. Y.; Murphy's, Col.; Red Bluff, Mont.

40.—KAOLIN.

Occurs in beds; it is a fine, white clay, plastic when wet; when dry is scaly or compact; can be crumbled in the fingers and feels gritty; adheres to the tongue; does not dissolve in acids.

VALUE.—Used for the finest porcelain and for adulterating candy.

LOCALITIES.—Found generally with iron-ore and fire-clay. Common on the eastern slope of the Alleghanies; Branford, Vt.; Beekman, Athol, Johnsburgh and McIntyre, N. Y.; Perth Amboy, N. J.; Reading, Tamaqua and New Gar-

den, Penn.; Mt. Savage, Md.; Richmond, Va.; Newcastle and Wilmington, Del.; Edgefield, S. C.; near Augusta, Ga.; Jacksonville, Ala.

#### 41.—LENTICULAR IRON ORE.

Occurs in beds or masses, consisting of minute flattened grains; little lustre; generally soils the fingers; breathed upon has a clayey odor; color, brownish-red, powder more red; dissolves in strong muriatic acid with some effervescence; brittle; gravity 4.

VALUE.—An ore of iron yielding thirty-three per cent. Generally mixed with other ores at the furnace.

LOCALITIES.—Found in sandstone. Wayne, Madison, Oneida and Herkimer Counties, N. Y.; Marietta O.

#### 42.—LEMONITE, OR BROWN HEMATITE.

Occurs in masses, with smooth rounded surfaces and fibrous structure; sometimes as hollow nodules, which are velvety-black inside; its powder when rubbed is yellowish-brown; when strongly heated turns

black; scratches glass feebly; brittle; dissolves in hot aqua-regia; gravity 4.

VALUE.—A common ore of pig-iron, containing sixty per cent.; used also for polishing buttons, etc.

LOCALITIES.—Found in heavy beds with mica-slate, quartz, limestone, etc. Salisbury and Kent, Conn.; Amenia, Fishkill, Dover and Beekman, N. Y.; Richmond and Lenox, Mass.; Pittsfield, Putney, Bennington and Ripton, Vt.; Hamburgh, N. J.; Pikeland and White Marsh, Penn.; Marquette, Mich.; Makequata River, Iowa; Iron Mountains, Stow and Green Counties, Mo.; Centerville, Ala.; near Raleigh and Smithfield, N. C.; on Coal Creek, Col.; and in coal areas generally.

#### 43.—MAGNETIC IRON ORE.

Occurs in granular masses, coarse or fine; attracted by the magnet, or affecting the compass-needle; powder black; brittle; dissolves in muriatic acid; gravity 5.

Series 10, Box 6, folder 3



Series 10, Box 6, folder 3

VALUE.—An important ore, yielding sixty-five per cent.

LOCALITIES.—Found in granite, slate and limestone rocks. Warren, Essex, Clinton, Saratoga, Herkimer, Orange and Putnam Counties, N. Y.; Raymond and Marshall's Island, Me.; Franconia, Jackson, Winchester, Lisbon, Swanzey and Unity, N. H.; Bridgewater, Chittenden, Marlboro, Rochester, Troy and Bethel, Vt.; Cambealon, R. I.; Hawley and Bernardston, Mass.; Haddam, Conn.; Goshen, Webb's Mine, Cornwall and White Marsh, Penn.; Hamburg, N. J.; Scott's Mills and Deer Creek, Md.; Mitchell and Madison Counties, N. C.; Spartanburg, S. C.; Laclede and Crawford Counties, Mo.; Sierra County, (Gold Valley,) Plumas, Tulare, Mariposa, Placer and El Dorado Counties, Cal.

44—MAGNETIC PYRITES.

Occurs massive; brittle; deep orange-yellow; powder grayish-black; metallic lustre; tarnishes easily; slightly attracts

the compass-needle; melts at a high heat, giving off sulphur-fumes; gravity 4.5.

VALUE.—Affords sulphur, copperas and nickel.


LOCALITIES.—Found in granite regions, often with copper and iron ores. Stafford, Corinth and Shrewsbury, Vt.; Trumbull and Monroe, Conn.; Port Henry, Diana and Orange County, N. Y.; Hurdstown, N. J.; Gap Mine, Lancaster County, Pa.; Ducktown Mines, Tenn.

45—MALACHITE.

Occurs in incrustations with smooth surface and fibrous; powder paler green than the mineral; brittle; by heat crackles and turns black; effervesces in acids; takes a fine polish, showing bands or rings; gravity 4.

VALUE.—Used for jewelry and inlaid work.

LOCALITIES.—Found in copper and lead mines. Cheshire, Conn.; Brunswick and Schuyler's Mines, N. J.; Morgantown, Cornwall, near Nicholson's Gap, Perki-



men and Phoenixville Lead Mines, Pa.; Petapasco Mines, Md.; Davidson County N. C.; Polk County, Tenn.; Left Hand River and Mineral Point, Wis.; Falls of St. Croix, Minn.; Jefferson County and Mine la Motte, Mo.; Calaveras County, Cal.; Big Williams' Fork, Ariz.; Wild Cat Cañon and near Virginia City, Mont.

46.—MANGANESE SPAR.

Occurs in masses; glassy lustre; color flesh or rose-red; becomes black on exposure; tough; melted with borax gives a violet-blue color; gravity 3.5.

VALUE.—Used in glaring stone-ware.

LOCALITIES.—Found in granite regions, often with iron-ore. Blue Hill Bay, Me.; Cumington, Warwick and Plainfield, Mass.; Irasburg and Coventry, Vt.; Winchester, and Hinsdale, N. H.; Cumberland, R. I.; Franklin and Hamburg, N. J.

47.—MARBLE.

Occurs coarse and fine granular; frequently veined or mottled; brittle; can be cut with a knife; takes a polish; efferves-

ces with acids; reduced to quicklime by heat; a gray variety contains stems and joints of worm-like fossils; gravity 2.5.

LOCALITIES.—Brandon, Rutland, Dorset, Shoreham, Pittsford, Middlebury, Fairhaven, Cavendish, Lowell, Troy and Sudbury, Vt.; West Stockbridge, Egremont, Great Barrington, Lanesboro, New Ashford, Sheffield, New Marlboro, Adams, Cheshire and Stoneham, Mass.; Clinton, Essex, Dutchess, Oneondaga, Putnam, St. Lawrence, Warren and Westchester, Counties, N. Y.; Smithfield, R. I.; New Haven, Milford, Conn.; near Philadelphia, N. J.; Texas and Hagerstown, Md.; Lancaster County, Pa.; Jefferson and Genevieve Counties, Mo.; Knox and Sevier Counties, Tenn.; Joliet, Ill.; Cherokee and Macon Counties, N. C.; Marquette, Mich.; near Deep River and on the Michigan and Menominee Rivers, Wis.

48.—MICA.

Occurs in masses, which can be split into very thin, elastic leaves; pearly lus-

62 UNDERGROUND TREASURES.

tre; at a high heat becomes opaque; gravity 3.

VALUE.—Used for doors of stoves, etc.

LOCALITIES.—Found in granite regions. Buckfield, Freeport and Oxford, Me.; Acworth, Grafton and Alstead, N. H.; Chesterfield, Barre, Mendon, South Royalston, Brimfield, Gosben and Russell, Mass.; Monroe, Haddam and Middletown, Conn.; Warwick, Edenville, Edwards, Monroe and Greenfield, N. Y.; Pennsbury, Thornbury, Unionville, Middletown and Chestnut Hill, Pa.; Jones' Falls, Md.

49.—MICACEOUS IRON ORE.

Resembles *Specular Iron Ore*, but consists of thin shining scales or leaves; powder dark red; a thin flake is translucent, showing red light; feels somewhat slippery.

VALUE.—Used as an ore of iron and for polishing.

LOCALITIES.—Hawley, Mass.; Piermont, N. H.; Ticonderoga, N. Y.; Warwick, Penn.; Loudon County, Va.

THE MINERALS TREATED. 63

50.—NITRE.

Occurs in thin crusts, delicate needles, or disseminated through the loose earth in caves; glossy lustre; brittle; cool, saline taste; crackles and burns brightly on live coals; a little harder than *Gypsum*.

VALUE.—Used in the manufacture of gunpowder, fulminating powders, nitric acid, etc.

LOCALITIES.—Marion County, Ky.; White County, Tenn.; near Rosiclare, Ill.; Silver Peak, Nev.

51.—OXYD OF MANGANESE.


Occurs in masses and little columns, often with small rounded surfaces; one ore is soft enough to be impressed by the nail, and soils; the other will scratch glass faintly; heated with borax, makes a violet glass; dissolves in hot muriatic acid, giving forth a yellowish-green gas; gravity 4 to 5.

VALUE.—Used for bleaching and for obtaining oxygen.

LOCALITIES.—Found in granite regions.

Series 10, Box 6, Folder 3





often with iron-ore. Brandon, Bennington, Monkton, Irasburg and Chittenden, Vt.; Hillsdale, Westmoreland and Westchester, N. H.; Plainfield, West Stockbridge and Conway, Mass.; Salisbury and Kent, Conn.; Montgomery County, Md.; Lake Superior Mining Region; Dubuque, Iowa; Deep Diggings, Mo.; Red Island, Cal.; Martinsburg, N. Y.

52.—PLATINUM.

Occurs in grains or lumps; metallic, silvery lustre; can be hammered out; heavier and harder than silver; not dissolved in nitric acid; gravity 17.

VALUE.—Nearly equal to *Gold*. Used for making chemical and philosophical apparatus, for coating copper, brass, etc.

LOCALITIES.—Found in river-gravel with *Gold*. Rutherford County, N. C.; Klamath region, Cape Blanco, on Salmon River, South Fork of Trinity, Butte, Honcut Cañon and Wood's Creeks, and on Middle Fork of American River, Cal.; at Gold Flat, Nev.

53.—RED COPPER ORE.

Occurs in crystals and masses; cochineal-red; powder brownish-red; nearly opaque; brittle; dissolves in nitric acid; heated on charcoal yields a globule of copper; gravity 6.

VALUE.—Affords copper, (sixty per cent.) and blue vitriol.

LOCALITIES.—Found in trap regions with other copper ores. Schuyler's, Somerville, New Brunswick and Flemington Mines, N. J.; Cornwall, Pa.; Ladenton, N. Y.; Lake Superior Region. Not abundant.

54.—RED HEMATITE.

Occurs in compact masses, with rounded surfaces or kidney-shaped; fibrous structure; color brownish-red to iron-black; but powder invariably red; when black, the lustre is somewhat metallic, otherwise dull; brittle; scratches glass with difficulty; dissolves slowly in strong muriatic acid; gravity 4.5 to 5.

VALUE.—An ore of iron, yielding from

thirty-six to fifty per cent. In powder, used as pigment and for polishing metals.

LOCALITIES.—Found usually in beds with granite or limestone. Aroostook County and Hodgdon, Me.; Antwerp, Ticonderoga, Crown Point and Gouverneur, N. Y.; Vernon, N. J.; West Whiteland, Pa.; Chatham and Orange Counties, N. C.; Marquette, Mich.; Shasta County, Cal. This mineral graduates into a soft, earthy variety, called *red ochre*, and into a compact, slaty variety, called *red chalk*, which has a clayey odor when breathed on.

55.—RED SILVER ORE.

Occurs in crystals and masses; metallic lustre; brittle; powder cochineal-red; easily cut; at a high heat yields a silver globule; the powder heated with potash turns black; gravity 6.

VALUE.—An ore of silver yielding sixty per cent.

LOCALITIES.—Found at gold and silver mines. Washoe and Austin, Nev.; Poor-man Lode, Idaho.

56.—RENSSELAERITE.

Occurs in masses; wax-like; a trifle harder than marble; when fresh can be scratched by the nail; soapy feel; takes a polish; cleavable; gravity 2.8.

VALUE.—Used as a marble and worked into inkstands, etc.


LOCALITIES.—Found with *steatite*, *serpentine*, limestone, etc. Antwerp, Canton, Fowler, De Kalb, Edwards, Russell and Gouverneur, N. Y.

57.—ROCK CRYSTAL.

Occurs in crystals and masses; transparent; glassy lustre; colorless; tough; brittle; not acted upon by acids or heat; electric by friction; gravity 2.5.

VALUE.—Cut for ornaments, lenses, etc.

LOCALITIES.—Common in sandstone, limestone and iron ore. Paris, Me.; Benton and Bartlett, N. H.; Sharon and Woodstock, Vt.; Pelham and Chesterfield, Mass.; Ellenville, Little Falls, Watervliet, Fairfield, Middleville, Fowler, Antwerp, Rossie, Lake George and Pal-



atine, N. Y.; Minnesota Mine, Lake Superior; Ouachita Spring, Ark.

58.—ROCK SALT.

Occurs in irregular beds or masses; brittle; saline taste; crackles in the fire.

LOCALITIES.—Found with *gypsum*, clay and sandstone. Washington County, Va.; Petit Anse, La.; Silver Peak, Nev.; Salmon River Mountains, Oregon.

59.—RUTILE.

Occurs in crystals generally; metallic lustre; powder pale brown; brittle; unchanged by heat or acids; if powdered and fused with potash, then dissolved in muriatic acid, the solution boiled with tin-foil assumes a beautiful violet color; gravity 4.

VALUE.—Used for coloring porcelain and artificial teeth.

LOCALITIES.—Found in granite and limestone rocks. Warren, Me.; Merrimack, and Warren, N. H.; Bristol, Putney and Waterbury, Vt.; Windsor, Shelburne, Barre, Conway and Leyden, Mass.;

Monroe, Conn.; Warwick, Edenville, Amity and Kingsbridge, N. Y.; Sudsbury, West Bradford, Parksburg, Concord and Newlin, Pa.; Newton, N. J.; Crowder's and Clubb Mountains, N. C.; Habersham and Lincoln Counties, Ga.; Magnet Cave, Ark.


60.—SERPENTINE.

Occurs in masses; feeble, resinous lustre; color oily green; powder whitish; often yellowish gray on the outside; can be cut easily; takes a fine polish; becomes reddish by heat; gravity 2.5—same as *Marble*.

VALUE.—Worked into mantels, jambs, table-tops, and many other ornaments.

LOCALITIES.—Found as a rock in large masses. Deer Isle, Me.; Baltimore, Cavendish, Jay and Troy, Vt.; Newbury, Blanford, Middlefield and Westfield, Mass.; Newport, R. I.; near New Haven and Milford, Conn.; Fort Henry, Antwerp, Syracuse, Warwick, Phillipstown, Canton, Gouverneur, Johnsburg, Davenport's Neck, New Rochelle and Rye, N. Y.;





Frankford, Hoboken and Montville, N. J.; Texas, Pa.; Cooptown, Md.; Patterson, N. C.; Calaveras County, Cal.; Alder Gulch, Mont. Marble veined with serpentine is called *verd-antique*.

61.—SILICATE OF COPPER.

Occurs in incrustations and masses; color bluish-green; not fibrous; surface smooth; easily cut; does not effervesce in acid; blackens by heat; gravity 2.

VALUE.—An ore of copper, yielding thirty per cent.

LOCALITIES.—Found with other copper ores. Somerville and Schuyler's, N. J.; Morgantown and Cornwall, Pa.; Wolcottville, Conn.; Big Williams' Fork, Ariz.

62.—SILVER.

Occurs in masses, or strings and threads penetrating rocks and native copper and *galena*; metallic lustre; tarnishes grayish black; can be cut in slices and hammered out; dissolved in muriatic acid, it turns black on exposure; gravity 10.

LOCALITIES.—Chiefly found with cop-

per near trap-rocks, and in fine grained *galena* and dark brown *blende*. Gold contains from one to fifteen per cent. Bridgewater, N. J.; Davidson and Stanley Counties, N. C.; Lake Superior Region; Poorman's Lode, Idaho; Comstock Lode and Montezuma Ledge, Nev.; Alpine County and Maris Vein, Cal.; Clear Creek County, Col.

63.—SILVER GLANCE.

Occurs in small lumps, plates and threads; color dark gray; cuts like lead; melts in a candle giving off sulphur fumes; gravity 7.

VALUE.—The most important ore of silver, containing eighty-seven per cent.

LOCALITIES.—May be found almost everywhere, except in the coal regions; associated with other ores, quartz, limestone, baryta, etc. Most abundant where mineral veins cross one another. Comstock Lode, Gold Hill, Reese River, Cortez District and Silver-Sprout Vein, Nev.; Clear Creek County, Nev.

64.—SMALTINE.

Occurs in crystals and masses; metallic lustre; color tin-white to steel-gray; powder dark gray; brittle; gives off garlic odor in a candle; melted with borax makes a deep blue glass; gravity 6.5 to 7.

VALUE.—An ore of cobalt and arsenic, containing eighteen to seventy per cent.

LOCALITIES.—Found in veins in granite regions with other ores, Mine la Motte, Mo.; Chatham, Conn.

65.—SMITHSONITE.

Occurs in masses, often rounded, covered with minute crystals, or honey-combed; color white, dirty yellow or stone color; glassy lustre; brittle; effervesces in nitric acid; barely scratches glass; barely translucent; gravity 4.4.

VALUE.—Yields fifty per cent. of zinc.

LOCALITIES.—Found generally in limestone with *galena* and *blende*. Friedenville, Lancaster and Perkiomen, Pa.; Linden and Mineral Points, Wis.; Lawrence, County, Ark.; Ewing's Diggings, Minn.

Series 10, Box 6, Folder 3

66.—SPATHIC IRON.

Occurs in crystals or plates somewhat curving; also (in coal regions) in nodules with concentric layers like an onion; brittle; color varies from white to yellowish-brown or dark-brown; strongly heated it blackens and will then attract the compass needle; the powder effervesces in nitric acid; melted with borax makes a green or yellow glass; gravity 3.8.

VALUE.—Yields thirty per cent. of iron, well adapted for steel.

LOCALITIES.—Found in granite and coal-formations, often with other ores. Plymouth, Vt.; Sterling, Mass.; Roxbury, Conn.; Antwerp, Herfman and Rossie, N. Y.; Fentress and Harlem Mines, N. C.; Coal Regions of Western Pa., Virginia, Eastern Ohio, etc.

67.—SPECULAR IRON ORE.

Occurs crystallized and in large masses, high metallic lustre; color steel-gray or iron-black; brittle; opaque except when very thin; the powder when very fine and

74 UNDERGROUND TREASURES.

rubbed on white paper shows red; the powder dissolves slowly in muriatic acid; by a strong heat yields a black mass which attracts the needle; gravity 5.

VALUE.—Yields from fifty to seventy per cent. of iron.

LOCALITIES.—Found in granite regions. Marquette, Mich.; Pilot Knob and Iron Mountains, Mo.; St. Lawrence County, N. Y.; Bartlett, Lisbon and Franconia, N. H.; Chittenden and Weathersfield, Vt.; Sauk County, Wis.

68.—SPINEL RUBY.

Occurs in pyramidal crystals; glassy lustre; powder white; scratches *rock-crystal*; by heat becomes black; gravity 3.5.

VALUE.—A gem; clear specimens weighing over four carats, are valued at half the price of the *diamond*.

LOCALITIES.—Found in granular limestone and clay. Amity and Gouverneur, N. Y.; Franklin and Byram, N. J.; Bolton and Boxborough, Mass.

10 BOX 6  
Series 10, Box 6, Folder 3  
THE MINERALS TREATED. 75

69.—STEATITE.

Occurs in masses, consisting of minute pearly scales or grains; can be marked by the nail; hardens by heat; soapy feel; gravity 2.5.

VALUE.—Used for fire-stones, tubes, in manufacture of porcelain, etc.

LOCALITIES.—Found in beds with limestone, serpentine and slate. Orr's Island, Me.; Frankestown, Keene, Orford and Pelham, N. H.; Athens, Cavendish, Marlboro, Moreton, New Fane, Bradboro, Troy, Waterville, Westfield, Weathersfield and Windham, Vt.; Middlefield, Lenox and Westfield, Mass.; Manayunk and Chestnut Hill, Pa.; Albemarle and Loudon Counties, Va.; Staten Island and St. Lawrence County, N. Y.; Bare Hills, Md.

70.—STRONTIANITE.

Occurs in crystals and in fibrous or granular masses; glassy lustre; brittle; thin pieces melt before a blow-pipe tinging the flame red; effervesces with acids; gravity 3.6.



VALUE.—A source of nitrate of strontia used in fire-works.

LOCALITIES.—Found in limestone. Schoharie, Muscalonge Lake, Chaumont Bay and Theresa, N. Y.

71.—SULPHUR.

Occurs in crystals, masses and crusts; brittle; can be easily cut; burns with a blue flame and sulphur odor; gravity 2.

LOCALITIES.—Found in limestone and gypsum, and around geysers and sulphur springs. Springport, N. Y.; on the Potomac, twenty-five miles above Washington; Put-in-Bay Island, Lake Erie; Clear Lake, Cal.; Santa Barbara County, Col.; Humboldt County, Nev.

72.—TIN ORE.

Occurs in crystals, grains and masses; high lustre; powder gray or brownish; brittle; will strike fire with steel; unaltered by heat or acids; gravity 7,—being nearly as heavy as lead-ore.

VALUE.—The only ore of tin, containing seventy-nine per cent. No gold-mine ever

Series 10, Box 6, folder 3

paid such profits as the tin mines of Cornwall.

LOCALITIES.—Jackson, N. H.; Temescal, Cal.; Boonville, Idaho; near Fredericktown, Mo.

73.—TOPAZ.

Occurs in crystals; glassy lustre; brittle; scratches *rock-crystal*; not acted upon by ordinary heat or acids; gravity 3.5.

VALUE.—A gem; the most esteemed are the rose-red and white.

LOCALITIES.—Found in granite. Trumbull, Willimantic and Middletown, Conn.; Crowder's Mountain, N. C.; Thomas's Mountains, Utah.

74.—TOURMALINE.

Occurs in crystals, usually in long, slender three-sided prisms which break easily, glassy lustre; brittle; becomes milk-white by heat; scratches *rock-crystal* and *garnet*; gravity 3.

VALUE.—Used for jewelry.

LOCALITIES.—Found in granite rocks. Paris, Albany and Hebron, Me.; Chester-

78 UNDERGROUND TREASURES.

field and Goshen, Mass.; Newlin and Marple, Pa.

75.—VARIEGATED COPPER ORE.

Occurs in crystals and masses; metallic lustre; quickly tarnishes; color between copper-red and light-brown; powder pale grayish-black; dissolves in nitric acid; at a high heat melts to a copper globule; heated on charcoal gives off fumes of sulphur; gravity 5.

VALUE.—An important ore of copper yielding sixty per cent.

LOCALITIES.—Found in granite, freestone, etc., with other ores. Bristol and Cheshire, Conn.; Mahanoy, Pa.; Copper Mines of N. J.

76.—WAD.

Occurs in masses; earthy and loose; can be broken by the fingers, and soils; no lustre; melted with borax makes a violet glass; feels very light.

VALUE.—Used in bleaching and for making smalt.

Series 10, Box 6, Folder 3

THE MINERALS TREATED. 79

LOCALITIES.—Found in low places, generally in the vicinity of slate or iron ore beds. Warren, Vt.; Blue Hill, Hodgdon and Thomaston, Me.; Columbia and Duchess Counties, Austerlitz, Canaan Centre and Martinsburg, N. Y.; East Bradford and White Marsh, Pa.; Mine la Motte, Mo.

77.—WILLEMITE.

Occurs in crystals and masses; feeble lustre; brittle; can hardly be cut with a knife; sometimes scratches glass; makes a jelly in muriatic acid; gravity 4.


VALUE.—Contains seventy per cent. of zinc.

LOCALITIES.—Found in limestone with *sincite*. Franklin and Sterling, N. J.

78.—ZINCITE.

Occurs in foliated masses or grains, powder orange-yellow; brittle; dissolves in acids without effervescence; gravity 5.5.

VALUE.—Yields seventy-five per cent. of zinc.



**Names:**

Chapter IV -  
Prospecting Mineral

Riches



Series 10, Box 6, Folder 3

are traversed by metalliferous deposits; and, therefore, in exploring with a view to its mineral productions, no opportunity should be lost of observing the various sections thus naturally laid bare.

When fragments of an ore are found on a hill-side, it is very evident that the vein must lie higher up. If the vein is horizontal and the fragments are found on the top of the hill, there is no probability of finding much if any of the vein, for generally it has been washed away. Ore-veins, however, are almost always nearly vertical; so that boring is of little use, as it might pass by the richest vein, or, striking it lengthwise, give a too favorable result.

As heavy minerals do not drift far, metals are always found near their source.

Horizontal beds can be worked at the least cost.

Pockets and nodules, or any detached masses of minerals, are soon exhausted. Veins, lodes and beds are most valuable.

Boring a three-inch hole, which costs about \$1 a foot, is a good method of testing a mineral vein or bed which lies more or less horizontally. A shaft may be sunk in sandstone for from \$6 to \$3 per cubic yard; in slate and gravel, at from \$2 to \$1.

The existence of mineral springs, and the rapid melting of the snow in any locality, are no indications of ores.

SEARCHING FOR DIAMONDS. — Few things are so unpromising and unattractive as gems in their native state. Hence their slow discovery. There is little doubt that diamonds exist in many places as yet unknown, or where their presence is unsuspected. It is very difficult for the unpracticed eye to distinguish them from crystals of quartz or topaz. The color constitutes the main difficulty in detecting their presence. They are of various shades of yellowish brown, green, blue and rose-red, and thus closely resemble the common gravel by which they are surrounded.

Series 10, Box 6, folder 3

Often they are not unlike a lump of gum arabic, neither brilliant nor transparent. The finest, however, are colorless, and appear like rock-crystals.

In Brazil, where great numbers of diamonds, chiefly of small size, have been discovered, the method of searching for them is to wash the sand of certain rivers in a manner precisely similar to that employed in the gold fields, namely, by prospecting pans. A shovelful of earth is thrown into the pan, which is then immersed in water, and gently moved about. As the washing goes on, the pebbles, dirt and sand are removed, and the pan then contains about a pint of thin mud. Great caution is now observed, and ultimately there remains only a small quantity of sand. The diamonds and particles of gold, if present, sink to the bottom, being heavier, and are selected and removed by the practiced fingers of the operator. But how shall the gems be detected by one who has had no experience, and who in a jew-

eler's shop could not separate them from quartz or French paste? The difficulty can only be overcome by testing such stones as may be suspected to be precious. Let these be tried by the very sure operation of attempting to cut with their sharp corners glass, crystal or quartz. When too minute to be held between the finger and thumb, the specimens may be pressed into the end of a stick of hard wood and run along the surface of window glass. A diamond will make its mark, and cause, too, a ready fracture in the line over which it has traveled. It will also easily scratch rock-crystal, as no other crystal will.

But a more certain and peculiar characteristic of the diamond lies in the form of its crystals. The ruby and topaz will scratch quartz, but no mineral which will scratch quartz has the *curved edges* of the diamond. In small crystals this peculiarity can be seen only by means of a magnifying glass; but it is invariably present.

Series 10, Box 6, folder 3

Interrupted, convex or rounded angles, are sure indications of genuineness. Quartz crystal is surrounded by six faces; the diamond by four. The diamond breaks with difficulty; and hence a test sometimes used is to place the specimen between two hard bodies, as a couple of coins, and force them together with the hands. Such a pressure will crush a particle of quartz, but the diamond will only indent the metal.

The value of the diamond is estimated by the carat, which is equal to about four grains, and the value increases rapidly with its weight. If a small, rough diamond weigh four grains, its value is about \$10; if eight grains, \$40; if sixteen grains, \$640. A cut diamond of one carat is worth from \$50 to \$100.

The imperfections of the diamond, and, in fact, of all cut gems, are made visible by putting them into oil of cassia, when the slightest flaw will be seen.

A diamond weighing ten carats is "princely;" but one in ten thousand weighs so much.

If a rough diamond resemble a drop of clear spring water, in the middle of which you perceive a strong light; or if it has a rough coat, so that you can hardly see through it, but white, and as if made rough by art, yet clear of flaws or veins; or, if the coat be smooth and bright, with a tincture of green in it,—it is a good stone. If it has a milky cast, or a yellowish-green coat, beware of it. Rough diamonds with a greenish crust are the most limpid when cut.

Diamonds are found in loose pebbly earth, along with gold, a little way below the surface, towards the lower outlet of broad valleys, rather than upon the ridges of the adjoining hills.

SEARCHING FOR GOLD.—The paying localities of gold deposits are the slopes of the Rocky and Alleghany Mountains. Gold need not be looked for in the anthra-



Series 10, Box 6, folder 3

cite and bituminous coal-fields nor in limestone rock. It is seldom found in the beds of rivers. The thing itself is the surest indication of its existence. If soil or sand is "washed" as described in Chapter V., and the particles of gold are not heavy enough to remain at the bottom but float away, the bed will not pay.

Along streams rather high up among the mountains, and in the gravelly drift covering the slopes of the valley below, are the best prospects. Where the stream meets an obstacle in its path or makes a bend or has deep holes, there we may look for "pockets" of gold. Black or red sands are usually richest. Gold-bearing rock is a slate or granite abounding in rusty looking quartz veins, the latter containing iron pyrites or cavities. Almost all iron pyrites and silver ores, may be worked for gold. When the quartz veins are thin and numerous rather than massive, and lie near the surface, they are considered most profitable. Few veins can be worked with

profit very far down. As traces of gold may be found almost everywhere, no one should indulge in speculation before calculating the percentage and the cost of extraction. Gold-hunting, after all, is a lottery with more blanks than prizes.

The substances most frequently mistaken for gold are *iron pyrites*, *copper pyrites* and *mica*. The precious metal is easily distinguished from these by its malleability (flattening under the hammer) and its great weight, sinking rapidly in water.

SEARCHING FOR SILVER.—This metal is usually found with lead ore and native copper. Slates and sandstones intersected by igneous rocks as trap and porphyry, are good localities. Pure silver is often found in or near iron ores and the dark brown zinc blende. The Colorado silver lodes are porous at the surface and colored more or less red or green. Any rock suspected of containing silver should be powdered and dissolved in nitric acid.

Series 10, Box 6, Folder 3


Pour off the liquid and add to it a solution of salt. If a white powder falls to the bottom which upon exposure turns black, there is silver in it. Silver mines increase in value as in depth, whereas gold diminishes as we descend.

**SEARCHING FOR COPPER.**—The copper ores, after exposure, or after being dipped in vinegar, are almost invariably green on the surface. They are most abundant near trap dykes. The pyrites is generally found in lead mines, and in granite and clay-slate. Copper very rarely occurs in the new formations, as along the Atlantic and Gulf borders, and in the Mississippi Valley south of Cairo.

**SEARCHING FOR LEAD.**—Lead is seldom discovered in the surface soil. It is also in vain to look for it in the coal region and along the coast. It must be sought in steep hills, in limestone and slate rocks. A surface cut by frequent ravines or covered by vegetation in lines, indicates mineral crevices. The galena from the slate

is said to contain more silver than that from the limestone. The purest specimens of galena are poorest in silver; the small veins are richest in the more precious metal. A lead vein is thickest in limestone, thinner in sandstone and thinnest in slate.

**SEARCHING FOR IRON.**—Any heavy mineral of a black, brown, red or yellow color may be suspected to be iron. To prove it, dissolve some in oil of vitriol and pour in an infusion of nut-gall or oak-bark; if it turns black, iron is present. If a ton of rich magnetic ore costs more than \$4 at the furnace, good hematite more than \$3, and poor ores more than \$1.50 or \$2, they are too expensive to pay, unless iron is unusually high. Deep mining for iron is not profitable. Generally speaking, a bed of good iron ore, a foot thick, will repay the cost of stripping it of soil, etc., twelve feet thick. Red and yellow earths, called ochres, contain iron. Magnetic ore is easily found by a compass.



**Names:**

Chapter V - Assay of  
Ores

**Types:**

book



Series 10, Box 6, folder 3

well, a bushel of rock often yielding half an ounce.

Iron occurs in large masses or beds; but the other metals are scattered in fragments through sand or soil, or exist in veins running through rocks.

WASHING FOR GOLD AND PLATINUM.—This operation, called "panning," is the oldest and simplest method of extracting the precious metals. At the present time, it furnishes to Russia nearly all the gold produced in that empire. It is based on the principle that substances of different weights may be separated by means of water,—the heaviest going to the bottom first. To examine the bank or bed of a river, suspected to contain gold, fill a milk-pan with the sands and carry it to a tub or pool of quiet water. Dip it under, stirring the mass with one hand or a stick. Then pour off the muddy water, fill with fresh water stirring again, and again pour off the light sand, clay, etc. Scales of gold will sink fast; mica flakes will take their time.

Repeat this process till all the fine particles are washed off; then allow just enough water to enter the pan as will cover the sand. By shaking the pan and gradually lowering the side by which it is held, the light sand will flow off, leaving in the corner a heap of coarse sand. Put in a small quantity of water and turn the pan around so as to create a gentle current, when the precious metal, if there be any, can be easily detected,—the gold by its bright lustre, the platinum by its lead color, and both by their malleability. Particles of gold are of uniform color and are either flat or rounded; while other yellow grains are angular. Holding the pan in the sunshine, secure any glittering glassy crystals, and test them for diamonds or rock-crystals. A magnet will remove any particles of magnetic iron-ore.

ASSAY OF GOLD ORE.—Gold may be found in quartz rock, in iron and copper pyrites, and in silver ores.

To ascertain if any gold is present in

Series 10, Box 6, Folder 3

96 UNDERGROUND TREASURES.

quartz, reduce the rock to powder and sift it. A certain quantity, say half a peck, is then washed as above described, till a manageable quantity of sand is left. If there is any show of gold, dry the mass and put it in a bowl or glass dish, and add an ounce of quicksilver, stirring the mixture well with a wooden rod. The quicksilver, which will unite with every particle of gold which may be there, is then poured off into a soft leather (chamois) bag. This is squeezed to remove superfluous quicksilver, and a pasty amalgam is left, which is put into an iron vessel and heated red hot. The yellow powder remaining is mixed with saltpetre and melted, when a button of pure gold will be found in the crucible. Quartz ores should yield \$6 to the ton in order to pay.

To test pyrites for gold, reduce a given quantity to powder and wash as before; then roast the residue at a red heat. Upon cooling, add quicksilver and treat as just

TESTING FOR GOLD. 97

described. Pyrites should yield \$1 of gold to the bushel of ore to be profitable.

Native silver often contains gold. To separate them, carefully flatten the alloy with a smooth hammer on an anvil, and then boil it in strong nitric acid in a glass flask for about ten minutes. Carefully pour off the acid into a vial, and wash the powder in the flask (which is fine gold) with water and dry. To the liquid in the vial add a solution of common salt. The white powder which falls should be removed, washed with water, and fused with powdered chalk or iron filings; a button of pure silver is the result.

Any substance supposed to be or to contain gold may be tested by dissolving it powdered in aqua regia and then pouring in a solution of copperas; if there is gold, the reddish-brown precipitate, by rubbing, assumes a bright metallic lustre.

To tell whether a globule of silver has any gold in it, put it on a white porcelain dish and moisten it with a drop of nitric

Series 10, Box 6, Folder 3

acid: if it is pure silver, it will dissolve and retain its white color; if mixed with gold, it will soon turn gray or black.

To test the purity of gold, rub some of it off on a hard black flint slate, and apply to the mark a drop of aqua fortis. If the gold is pure, the yellow streak remains unchanged, but if alloyed it partly disappears; if it is only an imitation of gold, it vanishes altogether.

A ready method of finding the amount of gold in a quartz rock with considerable accuracy, is by taking the specific gravity of the rock (well cleaned) as given on page 13. If the gravity is not over 2.7, it contains little or no gold. If it is 3, it very likely is gold-bearing, although pyrites may be present. But if it is over 5, it is undoubtedly auriferous, and if 12, it is very rich in gold.

It is generally considered that the sand of any river is worth working for the gold it contains, provided it will yield twenty-four grains to the hundred weight.

**ASSAY OF SILVER ORE.**—Pure silver is easily recognized. But lead and copper ores often contain a large percentage of the precious metal.

To detect silver in lead ore, dissolve the powdered ore in strong nitric acid; pour off the liquid and insert a piece of pure copper. If silver is present, it will go to the bottom. Or, add to the liquid a solution of common salt, and it will instantly become cloudy or white. If lead ore yields three ounces of silver to a ton, it may be worked for the silver as well as the lead. In Colorado, the average value of silver-bearing galena is \$100 per ton.

To test the copper ores for silver, dissolve them in nitric acid; then add a few drops of muriatic acid, and if silver is present, a white curdy precipitate will fall to the bottom. Native copper, when polished, often shows white spots of silver.

To estimate the proportion of silver in lead ore, reduce a known quantity of the clear ore to powder, mix with a little dry



Series 10, Box 6, Folder 3

100 UNDERGROUND TREASURES.

soda and a few nails, and heat in a round-bottomed iron pot or crucible. The lead which is obtained should then be put in a cup having ashes at the bottom, and strongly heated in an open furnace. A globule of silver will be left, if any is present, and being weighed, the percentage can be found.

Rich silver ores may be reduced by mixing them with ten parts of common salt, and exposing the mass for hours in an open furnace, stirring it frequently. When cold reduce to powder and mix with an equal quantity of quicksilver and enough water to make a paste, and agitate the mixture for two days, when the amalgam will fall to the bottom. The amalgam is then squeezed in a leather bag and washed.

Silver glance will yield its metal by heating it before a blow-pipe.

ASSAY OF COPPER ORE.—When the ore is native copper and rock, as at Lake Superior, it should be pounded and the earthy matter washed away. Then mix

ASSAY OF COPPER. 101

with a little potash or soda and bring to a high heat in a crucible.

Other copper ores may be tested by dissolving them powdered in dilute aqua regia. The presence of silver will be shown by a white powder on the bottom. Then add considerable ammonia. If there is any copper a blue liquor will be produced. Strain this through tissue paper, and evaporate to dryness. Dissolve the residue in muriatic acid, and by putting in a piece of iron or zinc, the copper will fall down. Or, add to this solution pure potash; dry and weigh the powder thrown down; every 5 parts of it contains 4 parts of copper.

Gray copper and red copper ores may be assayed by heating with charcoal, (both powdered,) in a furnace. Malachite and azurite should be smelted with borax; Copper pyrites and silicate of copper with soda or powdered marble.

A ton of copper ore which contains ten per cent. of metal, pays \$25 at the furnace. The ore of copper when roasted, turns

black; and when thrown into nitric acid makes a sky-blue solution. A clean knife-blade put into this solution will be coated with copper.

**ASSAY OF IRON ORE.**—Take a known quantity of the ore in fine powder and mix thoroughly with dry borax (or with one part of fluor spar, one of charcoal and four of salt,) and expose it for an hour in a covered crucible lined with charcoal to a white heat in a wind-furnace for an hour. A button of iron will be found at the bottom, which determines the percentage.

**ASSAY OF ZINC ORE.**—If the weighed ore is roasted with powdered charcoal, white flowers of zinc will be formed on a piece of cold iron held over it. After thorough roasting, the residue should be weighed; the loss is the oxide of zinc, and every 100 parts of this contain 81 of metal.

All the ores of zinc will dissolve in either nitric or hot sulphuric acid.


**ASSAY OF TIN ORE.**—Tin-stone will

Series 10, Box 6, folder 3

yield up its metal if mixed with charcoal, borax and soda, and heated on the hearth of a furnace or before a blow-pipe.

The presence of tin may be tested by dissolving the metal thus roasted out, in aqua regia and adding a decoction of Brazil-wood: if the metal was tin, the liquid will be colored a beautiful crimson.

**ASSAY OF LEAD ORE.**—Both galena and cerussite are rich ores, and when abundant pay well. They are easily reduced by heat, the former being usually mixed with charcoal and iron filings. If a western backwoodsman wants shot or bullets, he kindles a fire in a hollow tree or an old stump, puts some galena on the charred wood, and melts it down. After cooling, he finds the metal at the bottom. The smelting of a ton of lead costs about \$6. The average price per ton of galena is \$30. When galena is dissolved in warm nitric acid, a clean plate of zinc placed in it will be coated with brilliant blades of lead; if the galena contains silver, a plate




**Names:**

Chapter VI - Mineral  
Springs

**Types:**

book





108 UNDERGROUND TREASURES.

are often only saline, carbonated or magnesia waters.

The most famous iron springs are at Saratoga, Sandlake and Catskill, N. Y.; West Bethel, Fryeburg, Eberne and Bethel, Me.; Schooley's Mountain in Washington, N. J.; Bedford, Pittsburg, Frankfort and York, Pa.; Brandywine Springs, Del.; Red Sweet Springs in Monroe County, Rawley's Spring in Rockingham County, and Huguenot Springs in Powhattan County, Va.; in Bath County, Ky.; Yellow Springs, O.; twenty miles east of Knoxville, Tenn.; Madison County, Geo.; Raymond and Lynchburg, Miss.; near Ogden City, Utah; near Mt. Shasta, Col.

**SULPHUR SPRINGS.**—These are easily recognized by their unpleasant odor, resembling that of rotten eggs. The water blackens silver and a solution of sugar of lead.

Sulphur springs are very numerous. The best known are at Saratoga, Sharon, Clifton, Avon, Manlius, Chittenango, Dry-

ALUM AND EPSOM SPRINGS. 109

den and Richfield, N. Y.; Highgate and Newburg, Vt.; Togus, Bethel and West Newfield, Me.; Shover's Valley, Carlisle and Doubling Gap, Pa.; Winchester and Warrenton, Va.; Greenbrier and Monroe Counties, W. Va.; Bath County, Ky.; White's Creek near Nashville and in Granger County, Tenn.; Spartanburg, S. C.; Butts County, Geo.; Tallahatta, Ala.; Tampa, Fla.; near Bitter Creek and Great Salt Lake, Utah; along the Yellowstone River, Mont.; Jackson, Cal.

**ACID OR ALUM SPRINGS.**—These waters have a more or less sour taste and redden blue litmus-paper.

They are found at Byron and Oak Orchard, N. Y.; Blossburg, Pa.; Bath, Richmond and Rockbridge, Va.

**MAGNESIAN OR EPSOM SPRINGS.**—These have a bitter taste. To test any water for magnesia, add to a glass of it a solution of phosphate of soda and some hartshorn; if magnesia is present, the liquid first be-

Series 10, Box 6, folder 3

comes turbid, and finally minute crystals fall to the bottom.

There are Epsom springs at Harrodsburg and Perryville, Ky.; Westport, O.; Raymond, Miss.; Orange County, Ind.; Scott County, W. Va.

**SALINE OR SALT SPRINGS.**—These contain a large percentage of common salt, and are recognized by their taste. They generally contain many ingredients, (generally seven or eight,) but the salt predominates. A well should contain at least ten per cent. of salt to pay for working. The Syracuse spring yields a bushel of salt to every thirty-three gallons; while the Great Salt Lake contains 22 per cent. Among the most important salt wells are those at Syracuse, Salina and Liverpool, N. Y.; Lubec, Me.; Shannondale, Va.; Bath County, Ky.; Athens County, O.; Hartford, Ind.; Saginaw, Mich.; Oneida, Idaho.

**THERMAL OR WARM SPRINGS.**—Any spring is so called, the temperature of

Series 10, Box 6, Folder 3

which throughout the year is above that of the soil around it. They generally occur near the line of junction between the granite or igneous rocks and the stratified rock (slate or limestone) resting upon its flanks. The temperature of such waters in the United States ranges from 73 to 200 degrees, the latter being reached by the Geysers of Montana. Many iron and sulphur springs are also thermal.

The most noted warm springs are at Lebanon, N. Y.; in Bath, Berkley, Monroe and Scott Counties, Va.; Buncombe Counties, N. C.; French Brood River, Tenn.; Meriwether County, Geo.; Washitaw, Ark.; Salt Lake Valley, Utah; near Pyramid Lake, Nev.; along the Malheur and Fall Rivers, Or.; Lincoln Valley, Idaho; on Gardiner's River, in Madison County, and especially in the Yellowstone Basin, Mont.

**ARTESIAN WELLS.**—To sink a flowing well with any reasonable prospect of success, it is essential that the spot selected




should be lower than land in the vicinity, although those higher elevations may be several miles away. The layers of the rocks, also, should dip *towards* the spot rather than away from it. The best indication, but not a certain one, is a *great* basin-shaped valley, to the centre of which the rocks dip on one or more sides. Sandy, lime and slate rocks are more propitious than granite.

OIL WELLS.—Where there are marks of disturbance and misplacement of the rocks, there the experienced sink wells. Rugged hills and sharply-defined valleys are, generally, signs of such dislocation. The line or "break" from which the rocks dip like the roof of a house is considered most favorable. There is no such thing as an "oil rock," for the oil is found at different depths, and the fissure containing it is more or less vertical. In Pennsylvania, the greatest flowing wells have been found in the third sand rock. No limestone has afforded any large supply of

Series 10, Box 6, Folder 3

oil. Coal in no large quantities is ever found upon or in the immediate vicinity of the oil territory. The "show of oil" increases in value as a sign, with the depth at which it is found. Especially is the finding of a large amount of imprisoned gas, though no oil may be present, regarded as a good indication that oil is near. In the bituminous coal region, a gas spring indicates the probable existence of oil in the rocks below. But generally, "surface shows" are seductive. The great oil belt runs south-westerly from Oil Creek, Pa., to Burning Springs, West Va. But Ohio, Kentucky, Tennessee, Georgia, Alabama, Missouri, Texas, Illinois, Indiana, Michigan and Southern California are also rich in petroleum.

Frances Cabaniss Roberts Collection: Series 10, Box 6, Folder 3  
Orton, James "Underground Treasures: How and Where to Find Them"  
Image 61 r10\_06-03-000-0061 [Contents](#) [Index](#) [About](#)



**Names:**

Chapter VII -  
Artificial Jewelry

**Types:**

book

Series 10, Box 6, Folder 3

116 UNDERGROUND TREASURES.

of calcined potash, 10 of calcined borax and one of arsenic, keeping the mixture melted for two or three days and then cooling very slowly. Each ingredient is separately reduced to a fine powder.

FALSE RUBY is made by fusing together of strass one ounce and six drams, glass of antimony 37 grains, and purple of cassius one grain; then add eight parts more of strass and fuse for thirty hours; cool and remelt pieces in a blow-pipe. Or, melt five ounces of strass and one dram of manganese.

FALSE TOPAZ can be made from 1008 grains of strass, 43 grains of glass of antimony and one grain of purple of cassius.

FALSE SAPPHIRE.—Add to eight ounces of strass 52 grains of pure oxide of cobalt.

FALSE EMERALD.—To one pound of strass add one dram of verdigris and fifteen grains of crocus martis. Or, take 2304 grains of strass, 21 grains of green oxide of copper, and one grain of oxide of chrome. Or, take an ounce and a half

FALSE CARNELIAN AND OPAL. 117

of rock-crystal, six drams of dry soda, two drams of dry borax, two drams of red lead, one dram of nitre, twenty grains of red oxide of iron, and ten grains of green carbonate of copper.

FALSE CARNELIAN.—Strass two pounds, glass of antimony one pound, rouge two ounces, manganese one dram.

FALSE AMETHYSTS AND OPALS are manufactured; but the fine opal defies imitation, and the amethyst is too common in nature to allow much margin for the "pastes."

In distinguishing true and false gems, no one character should be depended upon. All genuine stones will bear rough handling; if the merchant says "hands off," refuse to purchase. Any gem worth buying is worth testing.

First: try the *hardness*. The file will make no impression on the diamond and ruby, and will with difficulty scratch the other gems; while the "pastes" are easily marred. All the precious stones scratch



Series 10, Box 6, Folder 3

window glass, although opal will not attack common bottle glass. All imitations easily yield to sand. The sapphire is the hardest of colored gems, and opal is the softest. The emerald will hardly scratch rock-crystal; its counterfeit not at all. Topaz will scratch ordinary ruby, but will not touch sapphire.

Secondly: as to *weight*. This is the most accurate method, but the stone must be taken from its setting. The mode of taking the gravity has already been given (page 13), and the amount of each is stated in Chapter II. Garnet is the heaviest of gems; weighed in water it loses only one-fourth of its weight; *i. e.*, if a red garnet be suspended by a fine thread from a delicate balance and immersed in a glass of water under it, one-quarter of its ordinary weight in air must be added to the pan from which it is suspended to restore the equilibrium. In like manner, ruby and sapphire lose a little more. The diamond and white topaz lose two-sevenths

of their weight. Rock-crystal, amethyst, carnelian and agate lose five-thirteenths; and opal about one-half, being the lightest of gems. The emerald loses more than one-third.

As "paste" can be made so as to have the same specific gravity as the genuine article, this test alone can not be relied upon; but very few of the imitations are so carefully made. The test is very convenient in distinguishing gems of like color from each other, as oriental ruby, spinel ruby and red tourmaline, and green tourmaline and emerald.


Thirdly: characteristics depending on *light* and *electricity*. It is not easy to look through a diamond of the first water, while imitations readily permit objects to be seen through them. A very delicate and perfect test of a diamond, distinguishing it from all colorless gems, as white topaz, white sapphire and white zircon, but not from "pastes," is to look through it at a pin-hole in a card. This requires some dex-

Series 10, Box 6, folder 3


terity, and the gem should be fixed to a steady object by a bit of wax at a proper distance. A true diamond will show but one hole, all the others will show two. As white topaz, when large, is a magnificent stone, it is often palmed off for a diamond of great value; but this test is invariably certain.

A true diamond retains its brilliancy under water.


When a colored stone is placed in the path of the solar spectrum (the row of seven colors into which sunlight is separated by a prism), its color will vary with the portion of the spectrum which falls upon it; and two stones of the same color, but of a different nature, will exhibit different effects. Thus, a paste placed beside a fine colored gem, betrays its worthlessness. A simpler method of testing stones is to look at them through a bit of glass, colored red, yellow, blue or green. Every stone will exhibit, under this test, properties peculiar to itself, and by which its na-



SEARCHING FOR DIAMONDS








1st 2 pages missing

**Names:**

Chapter VIII - Gold  
in California

**Types:**

book



1st 2 pages missing

**Names:**

Chapter VIII - Gold  
in California

**Types:**

book

p. 130

(pp. 127-128 torn out, missing)

Series 10, Box 6, Folder 3

FIRST WASHING FOR GOLD. 131

to persuade some of his friends to go with him; but they thought it would be only a waste of time and money, so he went with Bennet for his sole companion.

He arrived at Coloma on the 7th of March, and found the work at the mill going on as if no gold existed in the neighborhood. The next day he took a pan and spade, and washed some of the dirt in the bottom of the mill-race in places where Marshall had found his specimens, and, in a few hours, Humphrey declared that these mines were far richer than any in Georgia. He now made a rocker and went to work washing gold industriously, and every day yielded to him an ounce or two of metal. The men at the mill made rockers for themselves, and all were soon busy in search of the yellow metal. Everything else was abandoned; the rumor of the discovery spread slowly. In the middle of March Pearson B. Reading, the owner of a large ranch at the head of the Sacramento valley, happened to visit Sutter's Fort, and



Series 10, Box 6, Folder 3

hearing of the mining at Coloma, he went thither to see it. He said that if similarity of formation could be taken as a proof, there must be gold-mines near his ranch; so, after observing the method of washing, he posted off, and in a few weeks he was at work on the bars of Clear Creek, nearly two hundred miles north-westward from Coloma. A few days after Reading had left, John Bidwell, now representative of the northern district of the State in the lower House of Congress, came to Coloma, and the result of his visit was that, in less than a month, he had a party of Indians from his ranch washing gold on the bars of Feather River, twenty-five miles north-westward from Coloma. Thus the mines were opened at far distant points.


The first printed notice of the discovery of gold, was given in the California newspaper published in San Francisco on the 15th of March. On the 29th of May the same paper, announcing that its publication would be suspended, says:—"The

whole country, from San Francisco to Los Angeles, and from the sea-shore to the base of the Sierra Nevada, resound with the sordid cry of *gold! gold! gold!* while the field is left half planted, the house half built, and everything neglected but the manufacture of picks and shovels, and the means of transportation to the spot where one man obtained one hundred and twenty-eight dollars' worth of the real stuff in one day's washing; and the average for all concerned, is twenty dollars per diem."

The first to commence quartz mining in California were Capt. Wm. Jackson and Mr. Eliason, both Virginians, and the first machine used was a Chilian mill.

The Reid Mine, in North Carolina, was the first gold mine discovered and worked in the United States, and the only one in North America from which, up to 1825, gold was sent to the Mint.

Frances Cabaniss Roberts Collection: Series 10, Box 6, Folder 3  
Orton, James "Underground Treasures: How and Where to Find Them"  
Image 70 r10\_06-03-000-0070 [Contents](#) [Index](#) [About](#)



**Names:**

Chapter IX - Silver in  
Nevada

**Types:**

book

Series 10, Box 6, folder 3

136 UNDERGROUND TREASURES.

1862 to 1865 inclusive, about \$48,000,000, of which about \$32,750,000 were silver and \$15,250,000 gold; the production for the year 1866 was little less than \$15,675,000.

In 1854, the total amount of silver produced, was, according to Professor Whitney, \$47,443,200, of which sum \$7,864,000 came from European, and \$39,451,200 from American mines. The produce of silver of the Comstock vein in the last three years was about twenty-three per cent. of the entire amount furnished by all the silver mines in the world. It exceeds the aggregate produce of all European countries, and equals that of the entire western coast of South America, which in 1854 was \$11,099,200: the only country to which it is inferior being Mexico.

Dr. Raymond, United States Mining Commissioner, estimates the bullion product for the year 1869 as follows:

|                       |              |
|-----------------------|--------------|
| California, . . . . . | \$20,000,000 |
| Nevada, . . . . .     | 14,000,000   |
| Montana, . . . . .    | 12,000,000   |


YIELD OF PRECIOUS METALS. 137

|                                  |             |
|----------------------------------|-------------|
| Idaho, . . . . .                 | \$7,000,000 |
| Colorado and Wyoming, . . . . .  | 4,000,000   |
| Oregon and Washington, . . . . . | 4,000,000   |
| Arizona, . . . . .               | 1,000,000   |
| New Mexico, . . . . .            | 500,000     |





Frances Cabaniss Roberts Collection: Series 10, Box 6, Folder 3  
Orton, James "Underground Treasures: How and Where to Find Them"  
Image 72 r10\_06-03-000-0072 [Contents](#) [Index](#) [About](#)



**Types:**  
book cover

**Frances Cabaniss Roberts Collection: Series 10, Box 6, Folder 3**  
**Orton, James "Underground Treasures: How and Where to Find Them"**

[Contents](#)   [Index](#)   [About](#)

**Table of Contents**

[Image 1](#) (r10\_06-03-000-0001)  
[Image 2](#) (r10\_06-03-000-0002)  
[Image 3](#) (r10\_06-03-000-0003)  
[Image 4](#) (r10\_06-03-000-0004)  
[Image 5](#) (r10\_06-03-000-0005)  
[Image 6](#) (r10\_06-03-000-0006)  
[Image 7](#) (r10\_06-03-000-0007)  
[Image 8](#) (r10\_06-03-000-0008)  
[Image 9](#) (r10\_06-03-000-0009)  
[Image 10](#) (r10\_06-03-000-0010)  
[Image 11](#) (r10\_06-03-000-0011)  
[Image 12](#) (r10\_06-03-000-0012)  
[Image 13](#) (r10\_06-03-000-0013)  
[Image 14](#) (r10\_06-03-000-0014)  
[Image 15](#) (r10\_06-03-000-0015)  
[Image 16](#) (r10\_06-03-000-0016)  
[Image 17](#) (r10\_06-03-000-0017)  
[Image 18](#) (r10\_06-03-000-0018)  
[Image 19](#) (r10\_06-03-000-0019)

[Image 20](#) (r10\_06-03-000-0020)  
[Image 21](#) (r10\_06-03-000-0021)  
[Image 22](#) (r10\_06-03-000-0022)  
[Image 23](#) (r10\_06-03-000-0023)  
[Image 24](#) (r10\_06-03-000-0024)  
[Image 25](#) (r10\_06-03-000-0025)  
[Image 26](#) (r10\_06-03-000-0026)  
[Image 27](#) (r10\_06-03-000-0027)  
[Image 28](#) (r10\_06-03-000-0028)  
[Image 29](#) (r10\_06-03-000-0029)  
[Image 30](#) (r10\_06-03-000-0030)  
[Image 31](#) (r10\_06-03-000-0031)  
[Image 32](#) (r10\_06-03-000-0032)  
[Image 33](#) (r10\_06-03-000-0033)  
[Image 34](#) (r10\_06-03-000-0034)  
[Image 35](#) (r10\_06-03-000-0035)  
[Image 36](#) (r10\_06-03-000-0036)  
[Image 37](#) (r10\_06-03-000-0037)  
[Image 38](#) (r10\_06-03-000-0038)

[Image 39](#) (r10\_06-03-000-0039)  
[Image 40](#) (r10\_06-03-000-0040)  
[Image 41](#) (r10\_06-03-000-0041)  
[Image 42](#) (r10\_06-03-000-0042)  
[Image 43](#) (r10\_06-03-000-0043)  
[Image 44](#) (r10\_06-03-000-0044)  
[Image 45](#) (r10\_06-03-000-0045)  
[Image 46](#) (r10\_06-03-000-0046)  
[Image 47](#) (r10\_06-03-000-0047)  
[Image 48](#) (r10\_06-03-000-0048)  
[Image 49](#) (r10\_06-03-000-0049)  
[Image 50](#) (r10\_06-03-000-0050)  
[Image 51](#) (r10\_06-03-000-0051)  
[Image 52](#) (r10\_06-03-000-0052)  
[Image 53](#) (r10\_06-03-000-0053)  
[Image 54](#) (r10\_06-03-000-0054)  
[Image 55](#) (r10\_06-03-000-0055)  
[Image 56](#) (r10\_06-03-000-0056)  
[Image 57](#) (r10\_06-03-000-0057)

[Image 58](#) (r10\_06-03-000-0058)  
[Image 59](#) (r10\_06-03-000-0059)  
[Image 60](#) (r10\_06-03-000-0060)  
[Image 61](#) (r10\_06-03-000-0061)  
[Image 62](#) (r10\_06-03-000-0062)  
[Image 63](#) (r10\_06-03-000-0063)  
[Image 64](#) (r10\_06-03-000-0064)  
[Image 65](#) (r10\_06-03-000-0065)  
[Image 66](#) (r10\_06-03-000-0066)  
[Image 67](#) (r10\_06-03-000-0067)  
[Image 68](#) (r10\_06-03-000-0068)  
[Image 69](#) (r10\_06-03-000-0069)  
[Image 70](#) (r10\_06-03-000-0070)  
[Image 71](#) (r10\_06-03-000-0071)  
[Image 72](#) (r10\_06-03-000-0072)  
[Table of Contents](#)  
[Name & Place Index](#)  
[About the Collection](#)

**Frances Cabaniss Roberts Collection: Series 10, Box 6, Folder 3**  
**Orton, James "Underground Treasures: How and Where to Find Them"**

[Contents](#)   [Index](#)   [About](#)

**Name & Place Index**

Chapter I - Introduction [8](#)

Chapter II - Directions - The Key [11](#)

Chapter III - Descriptive List of Minerals [14](#)

Chapter IV - Prospecting Mineral Riches [44](#)

Chapter IX - Silver in Nevada [70](#)

Chapter V - Assay of Ores [50](#)

Chapter VI - Mineral Springs [56](#)

Chapter VII - Artificial Jewelry [61](#)

Chapter VIII - Gold in California [66](#), [67](#)

Hartford, CT [3](#), [4](#)

Illustrations [7](#)

Orton, James [3](#), [4](#)

Raymond, Rossiter W. [5](#)

Scottboro, AL [2](#)

Snodgrass, John [2](#)

Table of Contents [7](#), [8](#)

Underground Treasures [1](#), [3](#), [4](#), [6](#)



# Frances Cabaniss Roberts Collection

**Preferred Citation:** Frances Cabaniss Roberts Collection, Archives and Special Collections, M. Louis Salmon Library, University of Alabama in Huntsville, Huntsville, AL.

**Collection Scope and Content:** The Collection of 114 Linear ft. includes a total of 156 Archival Boxes. The Frances Cabaniss Roberts collection covers the historical records of the Cabaniss Roberts family. This collection contains extensive correspondence records of the Cabaniss Roberts family circa 1830 to 1930.

**Archives/Special Collections Access Restrictions:** None

**Conditions Governing Use:** This material may be protected under U. S. Copyright Law (Title 17, U.S. Code) which governs the making of photocopies or reproductions of copyrighted materials. You may use the digitized material for private study, scholarship, or research. Though the University of Alabama in Huntsville Archives and Special Collections has physical ownership of the material in its collections, in some cases we may not own the copyright to the material. It is the patron's obligation to determine and satisfy copyright restrictions when publishing or otherwise distributing materials found in our collections.

**Provenance:** Gift of Johanna Shields on October 28, 2006.



THE UNIVERSITY OF  
ALABAMA IN HUNTSVILLE

**The UAH Archives and Special Collections  
M. Louis Salmon Library**